



## JEE Main – 2024

### 27<sup>th</sup> JANUARY 2024 (Evening Shift)

#### General Instructions

1. The test is of **3 hours** duration and the maximum marks is **300**.
2. The question paper consists of **3 Subjects** (Subject I: **Mathematics**, Subject II: **Physics**, Subject III: **Chemistry**). Each Part has **two** sections (Section 1 & Section 2).
3. **Section 1** contains **20 Multiple Choice Questions**. Each question has 4 choices (1), (2), (3) and (4), out of which **ONLY ONE CHOICE** is correct.
4. **Section 2** contains **10 Numerical Value Type Questions** Out of which **ONLY 5 (any)** questions have to be attempted. You will **NOT** be allowed to attempt the sixth question. If you wish to attempt any other question apart from the five already attempted, then you will have to delete any one response from the five previously answered and then proceed to answer the new one.  
The answer to each question should be **rounded off to the nearest integer**.
5. No candidate is allowed to carry any textual material, printed or written, bits of papers, pager, mobile phone, any electronic device, etc. inside the examination room/hall.
6. On completion of the test, the candidate must hand over the Answer Sheet to the **Invigilator** on duty in the Room/Hall. **However, the candidates are allowed to take away this Test Booklet with them.**

#### Marking Scheme

1. **Section – 1:** +4 for correct answer, –1 (negative marking) for incorrect answer, 0 for all other cases.
2. **Section – 2:** +4 for correct answer, –1 (negative marking) for incorrect answer, 0 for all other cases.

**SECTION-1**

This section contains 20 Multiple Choice Questions. Each question has 4 choices (1), (2), (3) and (4), out of which ONLY ONE CHOICE is correct.

- An urn contains 6 white and 9 black balls. Two successive draws of 4 balls are made without replacement. The probability, that the first draw gives all white balls and the second draw gives all black balls, is:  
 (1)  $\frac{3}{715}$  (2)  $\frac{5}{256}$  (3)  $\frac{5}{715}$  (4)  $\frac{3}{256}$
- If  $y = y(x)$  is the solution curve of the differential equation  $(x^2 - 4) dy - (y^2 - 3y) dx = 0$ ,  $x > 2$ ,  $y(4) = \frac{3}{2}$  and the slope of the curve is never zero, then the value of  $y(10)$  equals:  
 (1)  $\frac{3}{1+(8)^{1/4}}$  (2)  $\frac{3}{1-2\sqrt{2}}$  (3)  $\frac{3}{1+2\sqrt{2}}$  (4)  $\frac{3}{1-(8)^{1/4}}$
- The values of  $\alpha$ , for which  $\begin{vmatrix} 1 & \frac{3}{2} & \alpha + \frac{3}{2} \\ 1 & \frac{1}{3} & \alpha + \frac{1}{3} \\ 2\alpha + 3 & 3\alpha + 1 & 0 \end{vmatrix} = 0$ , lie in the interval.  
 (1)  $(-3, 0)$  (2)  $(-2, 1)$  (3)  $(0, 3)$  (4)  $\left(-\frac{3}{2}, \frac{3}{2}\right)$
- The position vectors of the vertices A, B and C of a triangle are  $2\hat{i} - 3\hat{j} + 3\hat{k}$ ,  $2\hat{i} + 2\hat{j} + 3\hat{k}$  and  $-\hat{i} + \hat{j} + 3\hat{k}$  respectively. Let  $l$  denotes the length of the angle bisector AD of  $\angle BAC$  where D is on the line segment BC, then  $2l^2$  equals:  
 (1) 50 (2) 49 (3) 42 (4) 45
- Let  $g(x) = 3f\left(\frac{x}{3}\right) + f(3-x)$  and  $f''(x) > 0$  for all  $x \in (0, 3)$ . If  $g$  is decreasing in  $(0, \alpha)$  and increasing in  $(\alpha, 3)$ , then  $8\alpha$  is:  
 (1) 18 (2) 24 (3) 0 (4) 20
- Let  $\alpha = \frac{(4!)!}{(4!)^3!}$  and  $\beta = \frac{(5!)!}{(5!)^4!}$ . Then:  
 (1)  $\alpha \in N$  and  $\beta \notin N$  (2)  $\alpha \notin N$  and  $\beta \in N$   
 (3)  $\alpha \in N$  and  $\beta \in N$  (4)  $\alpha \notin N$  and  $\beta \notin N$
- Consider the function  $f : (0, 2) \rightarrow R$  defined by  $f(x) = \frac{x}{2} + \frac{2}{x}$  and the function  $g(x)$  defined by  $g(x) = \begin{cases} \min\{f(t)\}, & 0 < t \leq x \text{ and } 0 < x \leq 1 \\ \frac{3}{2} + x, & 1 < x < 2 \end{cases}$ . Then,  
 (1)  $g$  is continuous but not differentiable at  $x = 1$   
 (2)  $g$  is not continuous for all  $x \in (0, 2)$   
 (3)  $g$  is continuous and differentiable for all  $x \in (0, 2)$   
 (4)  $g$  is neither continuous nor differentiable at  $x = 1$

8. Let the position vectors of vertices  $A$ ,  $B$  and  $C$  of a triangle be  $2\hat{i} + 2\hat{j} + \hat{k}$ ,  $\hat{i} + 2\hat{j} + 2\hat{k}$  and  $2\hat{i} + \hat{j} + 2\hat{k}$  respectively. Let  $l_1$ ,  $l_2$  and  $l_3$  be the lengths of perpendiculars drawn from the ortho centre of the triangle on the sides  $AB$ ,  $BC$  and  $CA$  respectively, then  $l_1^2 + l_2^2 + l_3^2$  equals:

(1)  $\frac{1}{3}$                       (2)  $\frac{1}{5}$                       (3)  $\frac{1}{4}$                       (4)  $\frac{1}{2}$

9. Let the image of the point  $(1, 0, 7)$  in the line  $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$  be the point  $(\alpha, \beta, \gamma)$ . Then which one of the following points lies on the line passing through  $(\alpha, \beta, \gamma)$  and making angles  $\frac{2\pi}{3}$  and  $\frac{3\pi}{4}$  with  $y$ -axis and  $z$ -axis respectively and an acute angle with  $x$ -axis?

(1)  $(1, 2, 1 - \sqrt{2})$                       (2)  $(3, -4, 3 + 2\sqrt{2})$   
(3)  $(3, 4, 3 - 2\sqrt{2})$                       (4)  $(1, -2, 1 + \sqrt{2})$

10. Let  $A$  and  $B$  be two finite sets with  $m$  and  $n$  elements respectively. The total number of subsets of the set  $A$  is 56 more than the total number of subsets of  $B$ . Then the distance of the point  $P(m, n)$  from the point  $Q(-2, -3)$  is:

(1) 8                      (2) 4                      (3) 10                      (4) 6

11. Let  $f: R - \left\{ \frac{-1}{2} \right\} \rightarrow R$  and  $g: R - \left\{ \frac{-5}{2} \right\} \rightarrow R$  be defined as  $f(x) = \frac{2x+3}{2x+1}$  and  $g(x) = \frac{|x|+1}{2x+5}$ . Then, the domain of the function  $f \circ g$  is:

(1)  $R - \left\{ -\frac{7}{4} \right\}$                       (2)  $R - \left\{ -\frac{5}{2} \right\}$                       (3)  $R - \left\{ -\frac{5}{2}, -\frac{7}{4} \right\}$                       (4)  $R$

12. Let  $R$  be the interior region between the lines  $3x - y + 1 = 0$  and  $x + 2y - 5 = 0$  containing the origin. The set of all values of  $a$ , for which the points  $(a^2, a+1)$  lie in  $R$ , is:

(1)  $(-3, -1) \cup \left( \frac{1}{3}, 1 \right)$                       (2)  $(-3, -1) \cup \left( -\frac{1}{3}, 1 \right)$   
(3)  $(-3, 0) \cup \left( \frac{2}{3}, 1 \right)$                       (4)  $(-3, 0) \cup \left( \frac{1}{3}, 1 \right)$

13. The integral  $\int \frac{(x^8 - x^2)dx}{(x^{12} + 3x^6 + 1)\tan^{-1}\left(x^3 + \frac{1}{x^3}\right)}$  is equal to:

(1)  $\log_e \left( \left| \tan^{-1} \left( x^3 + \frac{1}{x^3} \right) \right| \right) + C$                       (2)  $\log_e \left( \left| \tan^{-1} \left( x^3 + \frac{1}{x^3} \right) \right| \right)^{\frac{1}{3}} + C$   
(3)  $\log_e \left( \left| \tan^{-1} \left( x^3 + \frac{1}{x^3} \right) \right| \right)^{\frac{1}{2}} + C$                       (4)  $\log_e \left( \left| \tan^{-1} \left( x^3 + \frac{1}{x^3} \right) \right| \right)^3 + C$

- 14.** If  $2 \tan^2 \theta - 5 \sec \theta = 1$  has exactly 7 solutions in the interval  $\left[0, \frac{n\pi}{2}\right]$ , for the least value of  $n \in \mathbb{N}$ , then  $\sum_{k=1}^n \frac{k}{2^k}$  is equal to:
- (1)  $\frac{1}{2^{13}}(2^{14} - 15)$  (2)  $\frac{1}{2^{15}}(2^{14} - 14)$  (3)  $1 - \frac{15}{2^{13}}$  (4)  $\frac{1}{2^{14}}(2^{15} - 15)$
- 15.** If  $\alpha, \beta$  are the roots of the equation,  $x^2 - x - 1 = 0$  and  $S_n = 2023 \alpha^n + 2024 \beta^n$ , then:
- (1)  $2S_{11} = S_{12} + S_{10}$  (2)  $S_{11} = S_{10} + S_{12}$   
 (3)  $2S_{12} = S_{11} + S_{10}$  (4)  $S_{12} + S_{11} + S_{10}$
- 16.** Let  $e_1$  be the eccentricity of the hyperbola  $\frac{x^2}{16} - \frac{y^2}{9} = 1$  and  $e_2$  be the eccentricity of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, a > b$ , which passes through the foci of the hyperbola. If  $e_1 e_2 = 1$ , then the length of the chord of the ellipse parallel to the x-axis and passing through (0, 2) is:
- (1)  $4\sqrt{5}$  (2)  $\frac{10\sqrt{5}}{3}$  (3)  $\frac{8\sqrt{5}}{3}$  (4)  $3\sqrt{5}$
- 17.** For  $0 < a < 1$ , the value of the integral  $\int_0^\pi \frac{dx}{1 - 2a \cos x + a^2}$  is:
- (1)  $\frac{\pi^2}{\pi + a^2}$  (2)  $\frac{\pi}{1 + a^2}$  (3)  $\frac{\pi^2}{\pi - a^2}$  (4)  $\frac{\pi}{1 - a^2}$
- 18.** If  $\lim_{x \rightarrow 0} \frac{3 + \alpha \sin x + \beta \cos x + \log_e(1 - x)}{3 \tan^2 x} = \frac{1}{3}$ , then  $2\alpha - \beta$  is equal to:
- (1) 2 (2) 1 (3) 7 (4) 5
- 19.** The 20<sup>th</sup> term from the end of the progression  $20, 19\frac{1}{4}, 18\frac{1}{2}, 17\frac{3}{4}, \dots, -129\frac{1}{4}$  is:
- (1) -100 (2) -110 (3) -118 (4) -115
- 20.** Considering only the principal values of inverse trigonometric functions, the number of positive real values of  $x$  satisfying  $\tan^{-1}(x) + \tan^{-1}(2x) = \frac{\pi}{4}$  is:
- (1) 2 (2) more than 2 (3) 1 (4) 0

**SECTION-2**

**Section 2 contains 10 Numerical Value Type Questions Out of which ONLY 5 (any) questions have to be attempted.** The answer to each question should be rounded off to the nearest integer.

- 21.** If the area of the region  $\{(x, y) : 0 \leq y \leq \min\{2x, 6x - x^2\}\}$  is  $A$ , then  $12A$  is equal to \_\_\_\_\_.
- 22.** If the solution curve, of the differential equation  $\frac{dy}{dx} = \frac{x+y-2}{x-y}$  passing through the point  $(2, 1)$  is  $\tan^{-1}\left(\frac{y-1}{x-1}\right) - \frac{1}{\beta} \log_e \left( \alpha + \left(\frac{y-1}{x-1}\right)^2 \right) = \log_e |x-1|$ , then  $5\beta + \alpha$  is equal to:
- 23.** The coefficient of  $x^{2012}$  in the expansion of  $(1-x)^{2008} (1+x+x^2)^{2007}$  is equal to \_\_\_\_\_.
- 24.** The lines  $\frac{x-2}{2} = \frac{y}{-2} = \frac{z-7}{16}$  and  $\frac{x+3}{4} = \frac{y+2}{3} = \frac{z+2}{1}$  intersect at the point  $P$ . If the distance of  $P$  from the line  $\frac{x+1}{2} = \frac{y-1}{3} = \frac{z-1}{1}$  is  $l$ , then  $14l^2$  is equal to \_\_\_\_\_.
- 25.** Let the complex numbers  $\alpha$  and  $\frac{1}{\alpha}$  lie on the circles  $|z - z_0|^2 = 4$  and  $|z - z_0|^2 = 16$  respectively, where  $z_0 = 1 + i$ . Then, the value of  $100|\alpha|^2$  is \_\_\_\_\_.
- 26.** Let  $f(x) = \int_0^x g(t) \log_e \left( \frac{1-t}{1+t} \right) dt$ , where  $g$  is a continuous odd function.  
If  $\int_{-\pi/2}^{\pi/2} \left( f(x) + \frac{x^2 \cos x}{1+e^x} \right) dx = \left( \frac{\pi}{\alpha} \right)^2 - \alpha$ , then  $\alpha$  is equal to \_\_\_\_\_.
- 27.** Let  $A$  be a  $2 \times 2$  real matrix and  $I$  be the identity matrix of order 2. If the roots of the equation  $|A - xI| = 0$  be  $-1$  and  $3$ , then the sum of the diagonal elements of the matrix  $A^2$  is \_\_\_\_\_.
- 28.** If the sum of squares of all real values of  $\alpha$ , for which the lines  $2x - y + 3 = 0$ ,  $6x + 3y + 1 = 0$  and  $\alpha x + 2y - 2 = 0$  do not form a triangle is  $p$ , then the greatest integer less than or equal to  $p$  is \_\_\_\_\_.
- 29.** Consider a circle  $(x - \alpha)^2 + (y - \beta)^2 = 50$ , where  $\alpha, \beta > 0$ . If the circle touches the line  $y + x = 0$  at the point  $P$ , whose distance from the origin is  $4\sqrt{2}$ , then  $(\alpha + \beta)^2$  is equal to \_\_\_\_\_.
- 30.** The mean and standard deviation of 15 observations were found to be 12 and 3 respectively. On rechecking it was found that an observation was read as 10 in place of 12. If  $\mu$  and  $\sigma^2$  denote the mean and variance of the correct observations respectively, then  $15(\mu + \mu^2 + \sigma^2)$  is equal to \_\_\_\_\_.

**SUBJECT II: PHYSICS****MARKS: 100****SECTION-1**

This section contains 20 Multiple Choice Questions. Each question has 4 choices (1), (2), (3) and (4), out of which ONLY ONE CHOICE is correct.

31. An object is placed in a medium of refractive index 3. An electromagnetic wave of intensity  $6 \times 10^8 \text{ W/m}^2$  falls normally on the object and it is absorbed completely. The radiation pressure on the object would be (speed of light in free space  $3 \times 10^8 \text{ m/s}$ ):
- (1)  $2 \text{ Nm}^{-2}$       (2)  $6 \text{ Nm}^{-2}$       (3)  $36 \text{ Nm}^{-2}$       (4)  $18 \text{ Nm}^{-2}$
32. Given below are two statements: one is labelled as Assertion (A) and the other is labelled as Reason (R).  
**Assertion (A):** Work done by electric field on moving a positive charge on an equipotential surface is always zero.  
**Reason (R):** Electric lines of forces are always perpendicular to equipotential surfaces.  
 In the light of the above statements, choose the most appropriate answer from the options given below:
- (1) (A) is not correct but (R) is correct  
 (2) (A) is correct but (R) is not correct  
 (3) Both (A) and (R) are correct and (R) is the correct explanation of (A)  
 (4) Both (A) and (R) are correct but (R) is not the correct explanation of (A)
33. Primary side of a transformer is connected to 230 V, 50 Hz supply. Turns ratio of primary to secondary winding is 10 : 1. Load resistance connected to secondary side is  $46 \Omega$ . The power consumed in it is:
- (1) 10.0 W      (2) 11.5 W      (3) 12.0 W      (4) 12.5 W
34. During an adiabatic process, the pressure of a gas is found to be proportional to the cube of its absolute temperature. The ratio of  $\frac{C_p}{C_v}$  for the gas is:
- (1)  $\frac{7}{5}$       (2)  $\frac{3}{2}$       (3)  $\frac{9}{7}$       (4)  $\frac{5}{3}$
35. The threshold frequency of a metal with work function 6.63 eV is:
- (1)  $1.6 \times 10^{15} \text{ Hz}$       (2)  $16 \times 10^{12} \text{ Hz}$       (3)  $1.6 \times 10^{12} \text{ Hz}$       (4)  $16 \times 10^{15} \text{ Hz}$
36. Given below are two statements:  
**Statement (I):** The limiting force of static friction depends on the area of contact and independent of materials.  
**Statement (II):** The limiting forces of kinetic friction is independent of the area of contact and depends on materials.  
 In the light of the above statements, choose the most appropriate answer from the options given below:
- (1) Both Statement I and Statement II are correct  
 (2) Statement I is correct and Statement II is incorrect  
 (3) Both Statement I and Statement II are incorrect  
 (4) Statement I is incorrect and Statement II is correct

37. The atomic mass of  ${}_6\text{C}^{12}$  is 12.000000 u and that of  ${}_6\text{C}^{13}$  is 13.003354 u. The required energy to remove a neutron from  ${}_6\text{C}^{13}$ , if mass of neutron is 1.008665 u, will be:

(1) 4.95 MeV (2) 62.5 MeV (3) 49.5 MeV (4) 6.25 MeV

38. Given below are two statements: one is labelled as Assertion (A) and the other is labelled as Reason (R).

**Assertion (A):** The property of body, by virtue of which it tends to regain its original shape when the external force is removed, is Elasticity.

**Reason (R):** The restoring force depends upon the bonded inter atomic and inter molecular force of solid.

In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) Both (A) and (R) are true and (R) is the correct explanation of (A)  
 (2) Both (A) and (R) are true but (R) is not the correct explanation of (A)  
 (3) (A) is true but (R) is false  
 (4) (A) is false but (R) is true

39. Given below are two statements: one is labelled as Assertion (A) and the other is labelled as Reason (R).


**Assertion (A):** The angular speed of the moon in its orbit about the earth is more than the angular speed of the earth in its orbit about the sun.

**Reason (R):** The moon takes less time to move around the earth than the time taken by the earth to move around the sun.

In the light of the above statements, choose the correct answer from the options given below:


- (1) (A) is not correct but (R) is correct  
 (2) Both (A) and (R) are correct but (R) is not the correct explanation of (A)  
 (3) (A) is correct but (R) is not correct  
 (4) Both (A) and (R) are correct and (R) is the correct explanation of (A)

40. The truth table of the given circuit diagram is:



	A	B	Y		A	B	Y
(1)	0	0	0	(2)	0	0	0
	0	1	0		0	1	1
	1	0	0		1	0	1
	1	1	1		1	1	0
	A	B	Y		A	B	Y
(3)	0	0	1	(4)	0	0	1
	0	1	1		0	1	0
	1	0	1		1	0	0
	1	1	0		1	1	1

41. A heavy iron bar of weight 12 kg is having its one end on the ground and the other on the shoulder of a man. The rod makes an angle  $60^\circ$  with the horizontal, the weight experienced by the man is:  
 (1) 3 kg (2)  $6\sqrt{3}$  kg (3) 6 kg (4) 12 kg
42. When a polaroid sheet is rotated between two crossed polaroids then the transmitted light intensity will be maximum for a rotation of:  
 (1)  $45^\circ$  (2)  $30^\circ$  (3)  $60^\circ$  (4)  $90^\circ$
43. A bullet is fired into a fixed target loses one third of its velocity after travelling 4 cm. It penetrates further  $D \times 10^{-3}$  m before coming to rest. The value of  $D$  is:  
 (1) 3 (2) 2 (3) 4 (4) 5
44. A current of  $200 \mu\text{A}$  deflects the coil of a moving coil galvanometer through  $60^\circ$ . The current to cause deflection through  $\frac{\pi}{10}$  radian is:  
 (1)  $60 \mu\text{A}$  (2)  $120 \mu\text{A}$  (3)  $30 \mu\text{A}$  (4)  $180 \mu\text{A}$
45. Three voltmeters, all having different internal resistances are joined as shown in figure. When some potential difference is applied across A and B, their readings are  $V_1$ ,  $V_2$  and  $V_3$ . Choose the correct option.



- (1)  $V_1 = V_2$  (2)  $V_1 + V_2 = V_3$  (3)  $V_1 + V_2 > V_3$  (4)  $V_1 \neq V_3 - V_2$
46. The equation of state of a real gas is given by  $\left(P + \frac{a}{V^2}\right)(V - b) = RT$ , where  $P$ ,  $V$  and  $T$  are pressure, volume and temperature respectively and  $R$  is the universal gas constant. The dimensions of  $\frac{a}{b^2}$  is similar to that of:  
 (1)  $PV$  (2)  $R$  (3)  $RT$  (4)  $P$
47. Given below are two statements: one is labelled as Assertion (A) and the other is labelled as Reason (R).  
**Assertion (A):** In Vernier Calliper if positive zero error exists, then while taking measurements, the reading taken will be more than the actual reading.  
**Reason (R):** The zero error in Vernier Calliper might have happened due to manufacturing defect or due to rough handling.  
 In the light of the above statements, choose the correct answer from the options give below:  
 (1) Both (A) and (R) are correct and (R) is the correct explanation of (A)  
 (2) (A) is true but (R) is false  
 (3) (A) is false but (R) is true  
 (4) Both (A) and (R) are correct but (R) is not the correct explanation of (A)
48. A ball suspended by a thread swings in a vertical plane so that its magnitude of acceleration in the extreme position and lowest position are equal. The angle ( $\theta$ ) of thread deflection in the extreme position will be:

- (1)  $2 \tan^{-1}\left(\frac{1}{\sqrt{5}}\right)$  (2)  $2 \tan^{-1}\left(\frac{1}{2}\right)$  (3)  $\tan^{-1}\left(\frac{1}{2}\right)$  (4)  $\tan^{-1}(\sqrt{2})$



49. The total kinetic energy of 1 mole of oxygen at  $27^{\circ}\text{C}$  is:

[Use universal gas constant (R) = 8.31J/mole K]

- (1) 5670.5 J      (2) 5942.0 J      (3) 6845.5 J      (4) 6232.5 J

50. Wheatstone bridge principle is used to measure the specific resistance ( $S_1$ ) of given wire, having length

$L$ , radius  $r$ . If  $X$  is the resistance of wire, then specific resistance is;  $S_1 = X \left( \frac{\pi r^2}{L} \right)$ . If the length of the


wire gets doubled then the value of specific resistance will be:

- (1)  $\frac{S_1}{4}$       (2)  $S_1$       (3)  $\frac{S_1}{2}$       (4)  $2 S_1$

**SECTION-2**

**Section 2 contains 10 Numerical Value Type Questions Out of which ONLY 5 (any) questions have to be attempted.** The answer to each question should be rounded off to the nearest integer.

51. A parallel beam of monochromatic light of wavelength  $5000\text{\AA}$  is incident normally on a single narrow slit of width  $0.001\text{ mm}$ . The light is focused by convex lens on screen, placed on its focal plane. The first minima will be formed for the angle of diffraction of \_\_\_\_\_ (degree).
52. Two charges of  $-4\mu\text{C}$  and  $+4\mu\text{C}$  are placed at the points  $A(1, 0, 4)\text{m}$  and  $B(2, -1, 5)\text{m}$  located in an electric field  $\vec{E} = 0.20\hat{i}\text{ V/cm}$ . The magnitude of the torque acting on the dipole is  $8\sqrt{\alpha} \times 10^{-5}\text{ Nm}$ , where  $\alpha =$  \_\_\_\_\_.
53. A closed organ pipe  $150\text{ cm}$  long gives 7 beats per second with an open organ pipe of length  $350\text{ cm}$ , both vibrating in fundamental mode. The velocity of sound is \_\_\_\_\_  $\text{m/s}$ .
54. The magnetic field at the centre of a wire loop formed by two semicircular wires of radii  $R_1 = 2\pi\text{m}$  and  $R_2 = 4\pi\text{m}$ , carrying current  $I = 4\text{ A}$  as per figure given below is  $\alpha \times 10^{-7}\text{ T}$ . The value of  $\alpha$  is \_\_\_\_\_. (Centre O is common for all segments)







55. A series LCR circuit with  $L = \frac{100}{\pi}\text{ mH}$ ,  $C = \frac{10^{-3}}{\pi}\text{ F}$  and  $R = 10\Omega$ , is connected across an ac source of  $220\text{ V}$ ,  $50\text{ Hz}$  supply. The power factor of the circuit would be \_\_\_\_\_.
56. A body falling under gravity covers two points A and B separated by  $80\text{ m}$  in  $2\text{ s}$ . The distance of upper point A from the starting point is \_\_\_\_\_  $\text{m}$  (use  $g = 10\text{ ms}^{-2}$ ).
57. The radius of pressure metre attached with a closed pipe is  $4.5 \times 10^4\text{ N/m}^2$ . On opening the value, water starts flowing and the reading of pressure metre falls to  $2.0 \times 10^4\text{ N/m}^2$ . The velocity of water is found to be  $\sqrt{V}\text{ m/s}$ . The value of V is \_\_\_\_\_.
58. The electric potential at the surface of an atomic nucleus ( $z = 50$ ) of radius  $9 \times 10^{-13}\text{ cm}$  is \_\_\_\_\_  $\times 10^6\text{ V}$ .
59. If Rydberg's constant is  $R$ , the longest wavelength of radiation in Paschen series will be  $\frac{\alpha}{7R}$ , where  $\alpha =$  \_\_\_\_\_.
60. A ring and a solid sphere roll down the same inclined plane without slipping. They start from rest. The radii of both bodies are identical and the ratio of their kinetic energies is  $\frac{7}{x}$ , where  $x$  is \_\_\_\_\_.

**SECTION-1**

This section contains 20 Multiple Choice Questions. Each question has 4 choices (1), (2), (3) and (4), out of which ONLY ONE CHOICE is correct.





61. The final product A, formed in the following reaction sequence is:







62. Which of the following cannot function as an oxidizing agent?








63. Which among the following halide/s will not show  $\text{S}_{\text{N}}1$  reaction:





Choose the most appropriate answer from the options given below:



64. Identify B formed in the reaction.



65. The molecular formula of second homologue in the homologous series of mono carboxylic acid is:




66. Match List-I with List-II.

List-I (Reaction)		List-II (Reagent(s))
(A)		(I) $\text{Na}_2\text{Cr}_2\text{O}_7, \text{H}_2\text{SO}_4$
(B)		(II) (i) NaOH (ii) $\text{CH}_3\text{Cl}$
(C)		(III) (i) NaOH, $\text{CHCl}_3$ (ii) NaOH (iii) HCl
(D)		(IV) (i) NaOH (ii) $\text{CO}_2$ (iii) HCl
(1)	(A) – (IV), (B) – (I), (C) – (III), (D) – (II)	(2) (A) – (IV), (B) – (III), (C) – (I), (D) – (II)
(3)	(A) – (II), (B) – (I), (C) – (III), (D) – (IV)	(4) (A) – (II), (B) – (III), (C) – (I), (D) – (IV)

67. The quantity which changes with temperature is:

- (1) Molarity (2) Molality (3) Mole fraction (4) Mass percentage

68. The order of relative stability of the contributing structure is:



Choose the correct answer from the options given below:

- (1)  $\text{III} > \text{II} > \text{I}$  (2)  $\text{I} = \text{II} = \text{III}$  (3)  $\text{II} > \text{I} > \text{III}$  (4)  $\text{I} > \text{II} > \text{III}$

69. Identify the incorrect pair from the following:

- (1) Photography – AgBr (2) Haber process – Iron  
(3) Polythene preparation –  $\text{TiCl}_4, \text{Al}(\text{CH}_3)_3$  (4) Wacker process –  $\text{Pt Cl}_2$

70. The incorrect statement regarding conformations of ethane is:

- (1) Ethane has infinite number of conformations  
(2) The conformations of ethane are inter-convertible to one-another  
(3) The dihedral angle in staggered conformation is  $60^\circ$ .  
(4) Eclipsed conformation is the most stable conformation.

71. The technique used for purification of steam volatile water immiscible substances is:
- (1) distillation (2) steam distillation  
(3) fractional distillation (4) fractional distillation under reduced pressure

72. Which structure of protein remains intact after coagulation of egg white on boiling?
- (1) Primary (2) Secondary (3) Quaternary (4) Tertiary

73. Bond line formula of  $\text{HOCH}(\text{CN})_2$  is:



74. Identify from the following species in which  $d^2sp^3$  hybridization is shown by central atom:



75. Phenolic group can be identified by a positive:

- (1) Tollen's test (2) Lucas test  
(3) Phthalein dye test (4) Carbylamine test

76. Given below are two statements:

**Statement (I):** In the lanthanoids, the formation  $\text{Ce}^{+4}$  is favoured by its noble gas configuration.

**Statement (II):**  $\text{Ce}^{+4}$  is a strong oxidant reverting to the common +3 state.

In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) Both Statement I and Statement II are true  
(2) Statement I is true but Statement II is false  
(3) Statement I is false but Statement II is true  
(4) Both Statement I and Statement II are false

77. Which of the following statements is not correct about rusting of iron?

- (1) When pH lies above 9 or 10, rusting of iron does not take place.  
(2) Rusting of iron is envisaged as setting up of electrochemical cell on the surface of iron object.  
(3) Coating of iron surface by tin prevents rusting, even if the tin coating is peeling off.  
(4) Dissolved acidic oxides  $\text{SO}_2$ ,  $\text{NO}_2$  in water act as catalyst in the process of rusting.

78. Given below are two statements:






**Statement (I):** Oxygen being the first member of group 16 exhibits only – 2 oxidation state.

**Statement (II):** Down the group 16 stability of +4 oxidation state decreases and +6 oxidation state increases.





In the light of the above statements, choose the most appropriate answer from the options given below:

- (1) Both Statement I and Statement II are correct  
(2) Both Statement I and Statement II are incorrect  
(3) Statement I is correct but Statement II is incorrect  
(4) Statement I is incorrect but Statement II is correct

79. Major product formed in the following reaction is a mixture of:



80. Choose the correct having all the elements with  $d^{10}$  electronic configuration from the following:






**SECTION-2**

Section 2 contains 10 Numerical Value Type Questions Out of which ONLY 5 (any) questions have to be attempted. The answer to each question should be rounded off to the nearest integer.

81. Time required for completion of 99.9% of a First order reaction is \_\_\_\_\_ times of half life ( $t_{1/2}$ ) of the reaction.

82. The spin only magnetic moment value of square planar complex  $[\text{Pt}(\text{NH}_3)_2\text{Cl}(\text{NH}_2\text{CH}_3)]\text{Cl}$  is \_\_\_\_\_ B.M. (Nearest integer)  
(Given atomic number for Pt = 78)


83. Total number of compounds with Chiral carbon atoms from following is \_\_\_\_\_ .



84. The hydrogen electrode is dipped in a solution of pH = 3 at 25°C. The potential of the electrode will be - \_\_\_\_\_  $\times 10^{-2}$  V.

$$\left( \frac{2.303 RT}{F} = 0.059 \text{ V} \right)$$


85. The number of non-polar molecules from the following is \_\_\_\_\_ .





86. For a certain thermochemical reaction  $\text{M} \rightarrow \text{N}$  at  $T = 400 \text{ K}$ ,  $\Delta H^\ominus = 77.2 \text{ kJ mol}^{-1}$ ,  $\Delta S = 122 \text{ JK}^{-1}$ , log equilibrium constant (log K) is - \_\_\_\_\_  $\times 10^{-1}$ .

87. Volume of 3 M NaOH (formula weight 40 g  $\text{mol}^{-1}$ ) which can be prepared from 84 g of NaOH is \_\_\_\_\_  $\times 10^{-1} \text{ dm}^3$ .

88. Total number of ions from the following with noble gas configuration is \_\_\_\_\_ .



89. 9.3 g of aniline is subjected to reaction with excess of acetic anhydride to prepare acetanilide. The mass of acetanilide produced if the reaction is 100% completed is \_\_\_\_\_  $\times 10^{-1} \text{ g}$ .



90. 1 mole of PbS is oxidized by "X" moles of  $\text{O}_3$  to get "Y" moles of  $\text{O}_2$ .  $X + Y =$  \_\_\_\_\_ .